

#2



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 12 MAR 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02001250.6

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 02001250.6
Demande no:

Anmeldetag:
Date of filing: 17.01.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Polymun Scientific
Immunbiologische Forschung GmbH
Nussdorfer Lände 11
1190 Wien
AUTRICHE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Anti-idiotypic antibody inducing HIV-1 neutralizing antibodies

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07K16/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR

ANTI-IDIOTYPIC ANTIBODY INDUCING HIV-1 NEUTRALIZING ANTIBODIES

TECHNICAL FIELD

5 The present invention is in the fields of immunology and vaccine development and relates to antibodies that mimic an epitope on gp41 of HIV-1. The invention further relates to applications of the antibodies and fragments thereof, including their use for eliciting HIV-1 neutralizing antibodies in mammalian hosts and to HIV-1 neutralizing antibodies elicited by said
10 antibodies or antibody fragments. The invention also relates to pharmaceutical compositions, in particular vaccines, containing such antibodies or antibody fragments for either active or passive immunization to inhibit or prevent HIV-1 infection in mammalian individuals.

15 BACKGROUND OF THE INVENTION

Anti-idiotypic antibodies (Ab2) are directed against the antigen binding site of other antibodies (Ab1). Epitopes of diverse pathogenic antigens may be mimicked functionally and structurally by those Ab2s. Thus, Ab2s are able to
20 induce anti-anti-idiotypic antibodies (Ab3) that are Ab1-like and have similar biological properties [1].

Different kinds of Ab2s are distinguished after immunization with Ab1 [2, 3]. The Ab2 α (Ab2-alpha) recognizes idiotypes at the distal position from Ab1
25 paratope and does not interfere with Ab1 binding to the nominal epitope. Ab2s beta (Ab2 β) are directed to the paratope of Ab1s and bear the internal image of the antigen. Ab2 β can compete with the nominal antigen for binding to the idiotypic and to antigen-specific xenogeneic antibodies. A third category, termed Ab2 γ (Ab2-gamma) recognizes idiotopes physically close to
30 the binding site but does not behave as a surrogate antigen [4].

Moreover, internal image anti-idiotypic antibodies may exhibit identical three-dimensional conformation or amino-acid sequence homology with the epitope. This has been shown for the reovirus system, where the amino acid sequence of the idiotypic had the same motif as the virus hemagglutinin [5]. Thus the virus-receptor binding on susceptible cells can be prevented.

Generally, the advantages of anti-idiotypic antibodies over other vaccines applying peptide antigens or inactivated viruses can be manifold [6]. They do not involve any viral components, they are safe but do not miss any specificity by their mimicry of single epitopes. They can be produced easily in large quantities for immunization. Ab2s can even induce an immune response against single neutralizing epitopes that may not be immunogenic on the native antigen. They thus can break immunogenic unresponsiveness to certain antigens as has been demonstrated in neonatal mice [7]. Active immunotherapy with Ab2-based vaccines have proven very potent to induce antigen-specific Ab3 in cancer patients against tumor cells [8-11].

Ab2s that display internal images of antigen (Ab2 β) have also induced protective immunity against infective pathogens [12].

HIV-1 infections may be accessible to Ab2 therapy because of the importance of neutralizing antibodies in the control of HIV-1 infections [13]. The possibility to induce neutralizing Ab3 in various systems including HIV-1 has been described [14].

Cross-reactivity of HIV-1 antigens with various normal human antigens suggests a potential advantage of single epitope Ab2 vaccines over multiple epitope vaccines [15]. Different approaches for the induction of a neutralizing humoral immune response have been made with anti-idiotypic antibodies. The most promising experiments were undertaken with anti-CD4 specific mAbs, which are able to induce HIV-1 neutralizing antibodies *in vitro* [16, 17]. The anti-CD4 mAbs Leu3a and IOT4A were also tested in clinical studies [18, 19]

and were shown to induce a gp120 cross-reacting immune response that inhibits gp120 binding to CD4. Ab2s against V3 loop peptides were also generated and used in different *in vitro* studies [20-22]. In different animal models they induced Ab3s with the ability of binding the autologous peptides as well as gp120, but could not neutralize HIV-1 in an *in vitro* experiment. Only one Ab2 has been described having the potential as a vaccine against HIV-1 [23].

In principle, Ab2 beta antibodies raised against antibodies neutralizing HIV-1 might have an enormous potential for vaccine design. A vaccine against HIV-1 must induce both, cellular as well as humoral immune responses. HIV-1 specific CTL-responses can be induced by various experimental vaccines such as DNA-vaccines [24] or chimeric influenza live viruses administered as nasal spray and expressing HIV-1 epitopes [25]. Reasonable experimental vaccines inducing neutralizing antibody responses are so far generally not available. The human monoclonal antibody 2F5 (produced by hybridoma cell line ECACC Accession No. 90091704) broadly and potently neutralizes primary HIV-1 isolates [26]. Thus, mAb 2F5 is an extraordinary interesting tool for the design of an anti-HIV-1 vaccine leading to properly induced humoral immune responses. It recognizes the highly conserved six amino acid core epitope sequence ELDKWA (SEQ ID NO:1) on the virus envelope ectodomain gp41 [27]. 2F5-like specific antibodies are only rarely found in sera of HIV-1 infected individuals. HIVIG (pooled sera of more than 70 HIV-1 positive donators) do not show significant 2F5-like specific binding to gp160 and/or gp41. The region on gp41 to which 2F5 binds is obviously cryptic to the human immune system during natural infection. Different mechanisms might contribute to its immuno-cryptic behavior. One explanation might be that the ~~human~~ complement factor H which is abundantly present in normal human serum has a binding domain in that region of gp41 thus masking it against immune recognition [28]. This mAb, although being a broad and potent entry ~~blocker~~ (neutralizers) of HIV-1 binds only very weakly to free virus and HIV-1 infected cells. [30]. It is therefore concluded that this fusogenic region on

- 4 -

gp41 is only exposed during the event of fusion of the HIV-1 with the host cell. That might on the one hand give an additional explanation for the cryptic nature of these neutralizing epitopes during natural infection. On the other hand HIV-1 isolates that are neutralization insensitive to this antibody have not been found so far.

In earlier studies the core epitope motif ELDKWA (SEQ. ID NO:1) of 2F5 was integrated into different antigenic formats. The presentation of the ELDKWA-motif on the haemagglutinin of influenza life virus was able to induce long lasting mucosal immune responses detected as 2F5-like specific IgA's in mouse faeces upon repeated immunization as a mucosal nasal spray [25]. However, quantities of IgA sufficient to prove *in vitro* neutralization potency could not be extracted from the faeces samples. Other forms of presenting the 2F5-core epitope for immunization such as fusion to the hepatitis B surface antigen expressed in yeast induced very high 2F5-like specific ELISA-titers in immunized BALB/c mice. However, the sera of those animals did not show any significant *in vitro* neutralization potency [36]. Peptide versions of the 2F5 epitope were also poorly immunogenic.

20 SHORT DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the reduction of binding capacity of 2F5 to the precoated epitope after preincubation of 2F5 with Ab2 3H6. Serial dilutions of 2F5 starting with 100 ng/mL were preincubated with 50 ng/mL or 500 ng/mL 3H6 or with the unspecific Ab2 6H8. After 1 hour the dilutions were allowed to react with GGGLELDKWASL (SEQ ID NO:13) at the microtiter plate and 2F5 was detected.

Fig. 2 shows a neutralization assay of 10 µg/mL 2F5 and 2G12 with increasing amounts of 3H6 and irrelevant mouse IgG.

Fig. 3 shows the binding of ascites and serum samples of 3H6 immunized mice to gp160.

Fig. 4 shows the binding of ascites and serum samples of 3H6 immunized mice to GGGLELDKWASL (SEQ ID NO.13)

5 Fig. 5 shows the result of a competition assay of 3H6 immunized mice sera with 2F5 in an ELISA binding study to precoated GGGLELDKWASL (SEQ ID NO:13)

DESCRIPTION OF THE INVENTION

10

It is an object of the invention to generate antigen binding components which are capable to induce neutralizing humoral immune responses against HIV-1. A further object of the invention is to provide novel antibodies or antibody fragments which are able to induce HIV-1 neutralizing antibodies.

15

The invention discloses murine antibodies and fragments thereof which are directed against an antibody of the gp41 antigen on HIV-1. In particular, the invention describes the generation of anti-idiotypic murine Ab2 β directed against the human monoclonal antibody 2F5 which is known to react with a
20 cryptic neutralizing epitope on gp41 of HIV-1. One of the Ab2's of the present invention, i.e. the murine monoclonal Ab2 designated 3H6 (deposited under the provisions of the Budapest Treaty on the Deposit of Microorganisms at PHLS Porton Down, Salisbury, UK; ECACC Acc.No. 01100279) blocks the binding of the human Ab1 2F5 to a synthetic epitope containing at least the
25 core epitope sequence ELDKWA (SEQ ID NO:1) as well as to gp160. 3H6 is also able to diminish the HIV-1 neutralizing potency of 2F5.

The present invention also relates to Fab-fragments derived from 3H6 antibodies, which were shown to induce neutralizing immune responses in the
30 sera of human and non-human mammals applying a simple prime/boost regimen of immunization. Thus, 3H6 Fab-fragments successfully mimic an

epitope that apparently lacks immunogenicity or is only poorly immunogenic on the native HIV-1 during natural infection.

An "antibody fragment" in the context of the present invention is an antigen-
5 binding part of the whole antibody which retains the binding specificity of the whole antibody molecule. The term encompasses, for example, Fab-, F(ab')₂-, Fv- and scFv-fragments. Antibody fragments can be obtained by conventional techniques well known in the art such as proteolytic digestion of antibodies, for example by papain digest, or through genetic engineering.

10

The antibodies and antibody fragments of the present invention can also be prepared by synthetic methods known in the art, such as peptide synthesis or recombinant DNA technology, once the structural nature of the antibody fragments has been analyzed by crystallization studies, sequencing methods
15 and/or other methods well known in the art.

In a further embodiment the present invention relates to an antibody or antibody fragment that is associated, e.g. fused or coupled, with a molecule that improves the immunogenic nature of the antibodies induced by the
20 antibody fragment upon administration to a mammalian individual. For instance, the antibody fragment may be coupled to a structure, for instance a cytokine such as, e.g., IL-4, which enhances the immunoreactivity of said mammal and preferably causes an increased B-cell response, which is of particular importance in the treatment of HIV. The present invention also
25 relates to entire Ab₂ antibodies, particularly to antibody 3H6, associated (e.g. coupled, fused or linked) with any such immunoresponse stimulating molecule.

In another embodiment, the invention relates to "humanized" antibodies and
30 antibody fragments. "Humanized" means that a non-human antibody or antibody fragment is linked with part of a human antibody. Usually it means that the complementarity determining regions (CDR) from the variable regions

of a non-human, e.g. rodent, antibody are combined with framework regions from variable regions of a human antibody, e.g. from a human IgG. In doing so the fragment containing the antigen binding site of the non-human antibody is inserted into a suitable human antibody. Methods for "humanizing" antibodies
5 are state of the art. For instance, an exon encoding the variable region containing the CDR of the non-human antibody may be spliced into the framework sequence of a suitable human antibody. Further, a Fab-fragment of the non-human antibody may be coupled with a suitable Fc-fragment of a suitable human antibody. Suitable human frameworks may be found by
10 homology searches in databases on the basis of homology to the non-human antibody.

"Humanized" antibody fragments according to the present invention may also be derived from synthetic phage libraries which use randomized combinations
15 of synthetic human antibody variable regions, by selecting on 2F5 antibodies as antigen.

In a still further embodiment the invention relates to chimeric antibodies. "Chimeric" antibodies are antibodies in which the whole of the variable
20 regions of a non-human, in particular rodent, antibody are expressed along with human constant regions of antibodies. This may provide antibodies with human effector functions. "Chimeric" antibodies may be created on the molecular level by fusing non-human exons encoding variable regions containing the antigen binding site with human exons encoding antibody
25 constant regions.

Sera of 3H6 induced immune responses show both specific binding inhibition patterns of 2F5 on the antigen as well as significant in vitro neutralization potency. Furthermore, the anti-2F5-Ab2 β 3H6, while reacting strongly with
30 the homologous Ab1, does not recognize isotype matched control antibodies demonstrating high specificity for the idiotype with no cross-reactivity to irrelevant Ig's.

- 8 -

Thus, the anti-idiotypic antibody 3H6 may be used, either directly as a vaccine to induce protective and neutralizing immune responses, or as a tool contributing to the design of an HIV-1 vaccine that also successfully induces
5 protective and neutralizing B-cell responses. Ab2s can be suitable for vaccination of HIV-1 infected newborns, whose immune system might be tolerant to stimulation with other antigens [37, 38].

At the first glance Ab2 3H6 represents a vaccine antigen capable to induce
10 neutralizing immune responses. Immunization with Ab2 3H6 shall avoid the induction of antibody-dependent enhancement of infection responses (ADE) observed in polyclonal sera of HIV-1 positive individuals [39-42]. The Ab1 2F5 was passively administered in gram-quantities to human volunteers. No signs of ADE have been observed in these human volunteers despite the fact that
15 complement activation was detected [29]. As Ab2 3H6 induces 2F5-like specificities in the sera of immunized animals it can be expected that ADE-phenomena observed with polyclonal sera of HIV-positive individuals [43] can be avoided upon vaccination with Ab2 3H6.

20 For the use as a pharmaceutical the antibodies and antibody fragments of the present invention may be part of a composition together with a pharmaceutically acceptable carrier. Thus, in a further embodiment the present invention relates to pharmaceutical compositions, particularly vaccines, which comprise the Ab2 antibodies or antibody fragments of the
25 present invention together with a suitable, optionally immunogenic, carrier and/or adjuvant.

In yet another embodiment the present invention relates to Ab3 antibodies (anti-anti-idiotypic antibodies) elicited by the present Ab2 antibodies or
30 antibody fragments of the 3H6 type, as well as to pharmaceutical compositions, including vaccines, which comprise Ab3 antibodies or antibody fragments directed against Ab2 3H6 and preferably directed against the

binding epitope of Ab2 3H6. In a most preferred embodiment, these Ab3 antibodies or antibody fragments are of a nature such that they neutralize HIV-1. Pharmaceutical compositions containing such an Ab3 antibody or fragment thereof are particularly useful for passive immunization against HIV-
5 1.

The antibodies or antibody fragments of the present invention may be administered in dosages of about 0.5 to about 10 µg/kg body weight, and preferably boosted in one or more, usually periodic, intervals. The
10 pharmaceutical compositions of the present invention containing either Ab2 3H6 antibodies or fragments thereof, or anti-3H6 Ab3 antibodies or fragments thereof may be administered prophylactically to prevent an individual from getting HIV-1 infected, as well as therapeutically to inhibit or stop progression of disease in an HIV-1 infected individual.

15

Example 1: Generation of 3H6 antibodies

Animals:

Female BALB/c mice, 8-10 weeks old, were obtained from Charles River (Sulzfeld, Germany).

20

Antibodies:

Immunoglobulin subclass standards and isotype-specific anti-mouse antisera were purchased from Sigma (St. Louis, MA). Monoclonal anti-HIV-1 antibody 2F5 was purified and used as antigen binding fragment (Fab') for immunization. Antibodies were digested with Mercuripapain (Sigma) and the
25 digest purified with Protein-A-Sepharose 4 FF-column.

Immunization of animals:

50 µg of the 2F5 Fab' were injected intraperitoneal once in complete Freund's adjuvant (CFA) and after three and six weeks again with incomplete Freund's adjuvant (IFA). Four days before spleen-ectomization the animals
30 were boosted for the last time. Spleenocytes were prepared on day 46 after the first immunization.

Generation of Hybridoma secreting mouse IgG:

The serum titer of mice was used as criteria for the development of immunocompetent spleenocytes. Only mice with an anti-2F5 titer of at least 1:100,000 were used for fusion. According to standard protocols the spleen cells were fused with X63-Ag-8.653 cells and plated in 96-well plates. HAT-selection was started the day after fusion and supernatants of growing wells were first tested for mouse IgG production using a mouse mAb as standard protein. Polyclonal goat anti-mouse- γ -chain antibody (Sigma) was diluted 1:500 in coating buffer (0.1 N $\text{Na}_2\text{CO}_3/\text{NaHCO}_3$ pH 9.6) and 100 μL per well were precoated onto microtiter plates (ELISA grade I; Nunc, Denmark). After each incubation step the plates were washed three times with PBS containing 0.1 % Tween 20 adjusted to pH 7.2. The standard mouse mAb was serially diluted, starting with 200 ng/mL (dilution buffer contained 1 % BSA in washing buffer). 50 μL per well of standard or serially diluted sample were allowed to bind to the precoated well for 1 h, and after washing the plates, we used 50 μL of 1:5,000 diluted goat anti-mouse IgG + IgM poly HRP80 (RDI Inc.) as second antibody. After 1 h incubation time, the plates were stained with OPD and read at 492 nm (reference wavelength 620 nm).

For the screening of antibodies against the 2F5 idiotype, the supernatants of 2170 HAT-resistant Hybridoma were tested for IgG production. 65 positive clones were found. Approx. 10 % of supernatants of the IgG positive clones were also reacting with antibody 2F5.

Example 2: Identification of neutralizing antibodies

Idiotype binding assay:

For initial screening of idiotype binding antibodies, 96-well microtiter plates were coated overnight at 4°C with 2F5-IgG1, 2F5-IgG3 or 2G12, another gp120 recognizing human anti-HIV-1 antibody [31]. Samples were diluted 2⁸ in dilution buffer starting with 200 ng/mL. The Ab2s were visualized as described for the mouse IgG ELISA.

The antibodies 3H6, 4F6, and 6F8 could be stabilized and were tested for their binding capacity in ELISA with different precoatings. The anti-idiotypic antibody 1G1, directed against a different neutralizing anti-HIV-1 antibody, namely 2G12 (ECACC Acc. No. 93091517; for details see WO 96/33219), served as control. Table 1 shows the relative binding properties of Ab2s with different idiotypes. 4F6 and 6F8 reacted with all antibodies and seemed to recognize an epitope at the constant region. The antibody 1G1 specifically binds to 2G12, while 3H6 recognizes both isotypes of 2F5 and it was negative for 2G12.

10 Table 1:

Ab2	2F5 IgG1	2F5 IgG3	2G12 IgG1
4F6	+	+	+
6F8	+	+	+
3H6	+	+	-
1G1	-	-	+

Idiotypic inhibition assay of Ab2 3H6:

Competitive binding assays were performed as previously described [32]. Different concentrations of Ab2 (50, 500 ng/mL) were preincubated with 1:2 dilutions of 2F5 starting with 100 ng/mL. After 1 hour the antibodies were allowed to react with the antigen on the plate. The solid phase was coated with the synthetic 2F5 epitope GGGLELDKWASL (SEQ. ID NO.13) at a concentration of 2 μ g/mL and the final binding of 2F5 was detected as described [33].

20

The ELISA competition assay was designated to describe the remaining binding capacity of 2F5 to the ELDKWA-epitope after preincubation with Ab2. Ab2 3H6 inhibits or prevents the binding of 2F5 to the epitope depending on the concentrations applied. Fig. 1 describes the decline of 2F5 binding to the GGGLELDKWASL (SEQ. ID NO.13) precoated plate with increasing concentrations of 3H6. Even 50 ng/mL 3H6 could reduce the binding properties

of 31 ng/mL 2F5 by more than one third (37 % reduction of the OD) and 500 ng/mL resulted in 83% reduction of OD while the Ab2 6F8 did not diminish the binding of 2F5 to the epitope. Since the binding affinity of 2F5 to the GST-(glutathion S-transferase) ELDKWA fusion-protein is $1.7 \times 10^7 \text{ M}^{-1}$ [33] the inhibition pattern of Ab2 3H6 was acceptable for further studies.

Similar results were obtained when replacing GGGLELDKWASL with other synthetic peptides containing the ELDKWA epitope or slightly modified variants thereof, the variants preferably being selected from the group consisting of ELDNWA (SEQ ID NO.2), ELNKWA (SEQ ID NO.3), LEELDKWA (SEQ ID NO.4), LEELDNWA (SEQ ID NO.5), LELNKWA (SEQ ID NO.6), ELDKWAS (SEQ ID NO.7), ELDNWA (SEQ ID NO.8), ELNKWAS (SEQ ID NO.9), LEELDKWAS (SEQ ID NO.10), LEELDNWAS (SEQ ID NO.11), LELNKWAS (SEQ ID NO.12).

Virus neutralization assay with 3H6:

Interaction of 2F5 with the different Ab2s was further investigated in a neutralization assay as described previously [26]. The assay was performed with HIV-1 strain NL4-3 infected AA-2 cells (NIAID, Bethesda, MD) as indicator cells using p24 antigen as virus replication marker. Serial dilutions of Ab2 antibodies (starting concentration of 250 $\mu\text{g/mL}$) were mixed with constant amounts of 2F5 or 2G12 as negative control (10 $\mu\text{g/mL}$ of each neutralizing antibody) and pre-incubation with virus before addition of AA-2 cells and further incubated for 5 days. Virus replication was measured in a p24-ELISA at the time point of termination of the assay [34]. The ratios of p24 antigen production in Ab-containing cultures (V_n) to p24 antigen production in control cultures (V_o), taking into account the input p24, were calculated and Ab concentrations ($\mu\text{g/mL}$) causing x% neutralization were plotted. Each test included a virus titration to determine the actual tissue culture infectious dose 50 (TCID_{50}) and the viral titer was determined by linear regression analysis. Tests were considered to be valid in case of viral titers between 10^2 - 10^3 TCID_{50} .

Fig. 2 shows the in vitro neutralization of HIV-1 NL4-3 after preincubation of Ab2 3H6 with 2F5 and 2G12. 2F5 and 2G12 alone neutralize 90 % of the virus at a concentration of 3 $\mu\text{g/mL}$ and 99 % of the virus at 12 $\mu\text{g/mL}$ and 11 $\mu\text{g/mL}$, respectively (data not shown). The 2F5 and 2G12 concentration was kept constant at 10 $\mu\text{g/mL}$. 50 μL of 2F5 or 2G12 were incubated with serial dilution of the Ab2s, 3H6, 4F6, 6F8 and only 3H6 was able to deplete neutralization of 2F5 (4F6 and 6F8 are not plotted). Fig. 1b describes the per cent neutralization representing the remaining neutralizing capacity of 10 $\mu\text{g/mL}$ 2F5 and 2G12 with increasing amounts of 3H6. 3.9 $\mu\text{g/mL}$ of 3H6 extinguish the 2F5 neutralization properties and 1.9 $\mu\text{g/mL}$ are able to reduce neutralizing capacity of 2F5 for 5 %. This means that 3H6 and 2F5 react in nearly equimolar ratio, since the molecular weight of the two corresponding antibodies may vary due to oligomerization or cleavage. 3H6 is not able to reduce neutralization of 2G12 at concentration as high as 250 $\mu\text{g/mL}$. Ab2 3H6 was able to reduce 2F5 neutralization for 95 % at 3.9 $\mu\text{g/mL}$.

Example 3: Induction of anti-anti-idiotypic antibodies (Ab3)

Induction of Ab3 in BALB/c mice:

After purification of Ab2 (3H6) by affinity chromatography it was digested with Mercuripapain (Sigma) and the digest purified with Protein-A-Sepharose 4 FF-column. Three BALB/c mice were injected with 5 μg of the 3H6 Fab' intraperitoneally and after three weeks again, both times with Freund's incomplete adjuvant. Another week later the mice were sacrificed and blood and ascites samples were collected. The samples were analyzed by ELISA for IgG titer as described above.

Three BALB/c mice were immunized with 5 $\mu\text{g/mL}$ 3H6 Fab' fragment FIA solution. Unfortunately, after the second immunization the animals developed ascites and had to be sacrificed without finishing the immunization scheme. Serum and ascites were shown to have total IgG titers up to 2 mg/ml. The results of IgG titers, binding capacity and competition assay are summarized in Table 2. The binding to GGGLELDKWASL (SEQ. ID NO.13) and gp160 in an

ELISA was illustrated in Figure 2 by plotting the cut-off dilution of the polyclonal serum and ascites. End point titers (cut-off) were calculated as the reciprocal of the highest dilution that resulted in an OD at least two times greater than the OD obtained with corresponding preimmune IgG.

5 Table 2:

Sample	IgG titer [$\mu\text{g/mL}$]	Cut-off titer binding to gp160	Cut-off titer binding to ELDKWA	50 % inhibition of 2F5/gp160 binding	Neutralizing capacity (IC ₅₀)
Serum 1	2271	1280	1280	1:60	-
Serum 2	3776	1280	1280	1:120	1:14
Serum 3	2603	1280	1280	1:120	1:12
Ascites 1	1127	320	320	1:15	-
Ascites 2	1958	640	640	1:15	-
Ascites 3	1442	320	320	1:15	-

Characterization of Ab3:

Specificity of Ab3:

Serum and ascites samples of the 3H6 immunized mice were assayed for
 10 binding to the synthetic epitope GGGLELDKWASL (SEQ. ID NO.13) (1 $\mu\text{g/mL}$
 in coating buffer) and recombinant gp160 (1 $\mu\text{g/mL}$ in coating buffer). The
 mouse antibodies were detected as described above. In another specificity
 test the serial dilutions of Ab3 samples were coincubated with a constant
 amount of 2F5 on a gp160 precoated plate. The 2F5 binding capacity at
 15 increasing dilutions of Ab3, starting with 1:10, was examined.

The 2F5 antibody was utilized at 25 ng/mL and a serial dilution of Ab3
 samples starting with 1:10 was used as competitor. Ab3 sera that most
 potently competed with 2F5 binding on the antigen also exhibited in vitro
 20 neutralization potency. Fig. 3 illustrates the reduction of 2F5 binding to the
 precoated gp160 in terms of competition with the Ab3 samples.

Syncytium inhibition Assay:

The neutralizing activity of Ab3 sera and ascites was assessed by inhibition of syncytium formation of HIV-1 strain RF_{NT}. Serial dilutions of serum, ascites or control antibody (2F5) were pre-incubated with virus (10^2 - 10^3 TCID₅₀/ml) for 5 1h at 37°C before addition of AA-2 cells. Starting dilutions of serum and ascites were 1:5 in culture medium. Cells were incubated for 5 days before assessment of syncytium formation. Experiments were performed with 4 replicates per dilution step. The presence of at least one syncytium per well was considered as indication for HIV-1 infection. The 50% inhibiting 10 concentration (IC₅₀) was calculated according to the method of Reed and Muench [35]. Experiments included a virus titration of the inoculum to confirm the infectious titer.

The neutralizing properties of serum and ascites were tested with HIV-1 15 laboratory strain RF_{NT}. Samples were diluted initially 1:5, sterilized by 0.2 μ filtration and used at a starting concentration of 1:10 in the AA-2 syncytia inhibition assay in quadruplicates. The virus-titer (TCID₅₀/ml) was determined to be $10^{2.75}$. The ascites samples and serum 1 did not show any antiviral effect. Sera 2 and 3 reached an IC50 at dilutions 1:14.1 and 1:11.9 20 respectively. Serum 1 didn't show neutralization as well as the ascites samples.

References:

1. Herlyn, D., *et al.*, *Anti-idiotypic cancer vaccines: pre-clinical and clinical studies*. In Vivo, 1991. 5(6): p. 615-23.
- 5 2. Jerne, N.K., J. Roland, and P.A. Cazenave, *Recurrent idiotopes and internal images*. Embo Journal, 1982. 1(2): p. 243-7.
3. Nisonoff, A., *Idiotypes: concepts and applications*. Journal of Immunology, 1991. 147(8): p. 2429-38.
4. Ertl, H.C. and C.A. Bona, *Criteria to define anti-idiotypic antibodies carrying the internal image of an antigen*. Vaccine, 1988. 6(2): p. 80-4.
- 10 5. Bruck, C., *et al.*, *Nucleic acid sequence of an internal image-bearing monoclonal anti-idiotypic and its comparison to the sequence of the external antigen*. Proceedings of the National Academy of Sciences of the United States of America, 1986. 83(17): p. 6578-82.
- 15 6. Kennedy, R.C. and R. Attanasio, *Concepts of idiotypic-based vaccines for hepatitis B virus and human immunodeficiency virus*. Canadian Journal Of Microbiology, 1990. 36(11): p. 811-6.
7. Hiernaux, J., C. Bona, and P.J. Baker, *Neonatal treatment with low doses of anti-idiotypic antibody leads to the expression of a silent clone*. Journal Of Experimental Medicine, 1981. 153(4): p. 1004-8.
- 20 8. Bhattacharya-Chatterjee, M. and K.A. Foon, *Anti-idiotypic antibody vaccine therapies of cancer*. Cancer Treatment And Research, 1998. 94: p. 51-68.
9. Herlyn, D., *et al.*, *Anti-idiotypic vaccine in colorectal cancer patients*. Hybridoma, 1993. 12(5): p. 515-20.
- 25 10. Pohl, C., *et al.*, *CD30-specific AB1-AB2-AB3 internal image antibody network: potential use as anti-idiotypic vaccine against Hodgkin's lymphoma*. International Journal Of Cancer, 1993. 54(3): p. 418-25.
11. Rajadhyaksha, M., Y.F. Yang, and Y.M. Thanavala, *Immunological evaluation of three generations of anti-idiotypic vaccine: study of B and T cell responses following priming with anti-idiotypic, anti-idiotypic peptide and its MAP structure*. Vaccine, 1995. 13(15): p. 1421-6.
- 30 12. Sikorska, H.M., *Therapeutic applications of anti-idiotypic antibodies*. Journal Of Biological Response Modifiers, 1988. 7(4): p. 327-58.
- 35 13. Attanasio, R. and R.C. Kennedy, *Idiotypic cascades associated with the CD4-HIV gp120 interaction: principles for idiotypic-based vaccines*. International Reviews Of Immunology, 1990. 7(1): p. 109-19.

14. Kang, C.Y., *et al.*, *Development of HIV/AIDS vaccine using chimeric gag-env virus-like particles*. Biological Chemistry, 1999. 380(3): p. 353-64.
15. del Guercio, P., *AIDS and immune dysfunction. Alternative etiologic mechanisms*. Contributions To Microbiology And Immunology, 1989. 11: p. 289-304.
16. Sutor, G.C., *et al.*, *Neutralization of HIV-1 by anti-idiotypes to monoclonal anti-CD4. Potential for idiotypic immunization against HIV*. Journal Of Immunology, 1992. 149(4): p. 1452-61.
17. Shearer, M.H., *et al.*, *Idiotypic cascades associated with the CD4-HIV glycoprotein 120 interaction: immunization with anti-idiotypic antibodies induces anti-anti-idiotypic responses with anti-CD4 specificity and in vitro neutralizing activity*. Aids Research and Human Retroviruses, 2000. 16(1): p. 77-86.
18. Wilks, D., *et al.*, *Anti-idiotypic responses to immunization with anti-Leu 3a in human immunodeficiency virus-seropositive individuals*. Journal of Infectious Diseases, 1991. 163(2): p. 389-92.
19. Deckert, P.M., *et al.*, *CD4-imitating human antibodies in HIV infection and anti-idiotypic vaccination*. Journal Of Immunology, 1996. 156(2): p. 826-33.
20. Boudet, F., J. Theze, and M. Zouali, *Anti-idiotypic antibodies to the third variable domain of gp120 induce an anti-HIV-1 antibody response in mice*. Virology, 1994. 200(1): p. 176-88.
21. Fung, M.S., *et al.*, *Monoclonal anti-idiotypic antibody mimicking the principal neutralization site in HIV-1 GP120 induces HIV-1 neutralizing antibodies in rabbits*. Journal Of Immunology, 1990. 145(7): p. 2199-206.
22. Connelly, R.J., *et al.*, *Dual specificity of a monoclonal anti-idiotypic antibody for HIV-1 neutralizing monoclonals 110.3 and 110.4 as well as the V3 loop of gp120*. Virology, 1994. 205(2): p. 554-7.
23. Sperlagh, M., *et al.*, *Polyclonal anti-idiotypic antibodies mimicking gp120 of HIV-1*. Viral Immunology, 1994. 7(2): p. 61-9.
24. Ayyavoo, V., *et al.*, *Immunogenicity of a novel DNA vaccine cassette expressing multiple human immunodeficiency virus (HIV-1) accessory genes*. Aids, 2000. 14(1): p. 1-9.
25. Ferko, B., *et al.*, *Chimeric influenza virus replicating predominantly in the murine upper respiratory tract induces local immune responses*

- against human immunodeficiency virus type 1 in the genital tract. Journal of Infectious Diseases, 1998. 178(5): p. 1359-68.*
26. Purtscher, M., et al., *A broadly neutralizing human monoclonal antibody against gp41 of human immunodeficiency virus type 1. Aids Research and Human Retroviruses, 1994. 10(12): p. 1651-8.*
 27. Muster, T., et al., *A conserved neutralizing epitope on gp41 of human immunodeficiency virus type 1. Journal Of Virology, 1993. 67(11): p. 6642-7.*
 28. Stoiber, H., et al., *Inhibition of HIV-1 infection in vitro by monoclonal antibodies to the complement receptor type 3 (CR3): an accessory role for CR3 during virus entry? Molecular Immunology, 1997. 34(12-13): p. 855-63.*
 29. Stiegler, G., et al., *A potent cross-clade neutralizing human monoclonal antibody against a novel epitope on gp41 of Human Immunodeficiency Virus Type 1. submitted, 2001.*
 30. Gorny, M.K. and S. Zolla-Pazner, *Recognition by human monoclonal antibodies of free and complexed peptides representing the prefusogenic and fusogenic forms of human immunodeficiency virus type 1 gp41. Journal of Virology, 2000. 74(13): p. 6186-92.*
 31. Kunert, R., F. Rüker, and H. Katinger, *Molecular characterization of five neutralizing anti-HIV type 1 antibodies: identification of nonconventional D segments in the human monoclonal antibodies 2G12 and 2F5. Aids Research And Human Retroviruses, 1998. 14(13): p. 1115-28.*
 32. Bruderer, U., et al., *Analyses of affinity distributions within polyclonal populations of antigen-specific antibodies. Evaluation of their accuracy in population detection using monoclonal antibodies. Journal of Immunological Methods, 1992. 151(1-2): p. 157-64.*
 33. Kunert, R., et al., *Stable recombinant expression of the anti HIV-1 monoclonal antibody 2F5 after IgG3/IgG1 subclass switch in CHO-cells. Biotechnology and Bioengineering, 2000. 67(1): p. 97-103.*
 34. Steindl, F., et al., *A simple and robust method for the complete dissociation of HIV-1 p24 and other antigens from immune complexes in serum and plasma samples. Journal of Immunological Methods, 1998. 217(1-2): p. 143-51.*
 35. Reed, L.J. and H. Muench, *A simple method of estimating fifty per cent endpoints. Am. J. Hygiene, 1938. 27: p. 493-497.*

36. Eckhart, L., *et al.*, *Immunogenic presentation of a conserved gp41 epitope of human immunodeficiency virus type 1 on recombinant surface antigen of hepatitis B virus*. Journal of General Virology, 1996. 77 (Pt 9): p. 2001-8.
- 5 37. Hanson, L.A., *et al.*, *Anti-idiotypic antibodies from the mother may influence the immune response of the fetus and newborn*. Acta Paediatrica Japonica, 1994. 36(2): p. 123-7.
38. Hanson, L.A., *et al.*, *Immunoregulation mother-fetus/newborn, a role for anti-idiotypic antibodies*. Acta Paediatrica Scandinavica. Supplement, 1989. 351: p. 38-41.
- 10 39. Forthal, D.N., *et al.*, *Functional activities of 20 human immunodeficiency virus type 1 (HIV-1)-specific human monoclonal antibodies*. Aids Research and Human Retroviruses, 1995. 11(9): p. 1095-9.
- 15 40. Kozlowski, P.A., *et al.*, *High prevalence of serum IgA HIV-1 infection-enhancing antibodies in HIV-infected persons. Masking by IgG*. Journal of Immunology, 1995. 154(11): p. 6163-73.
41. Mascola, J.R., *et al.*, *Summary report: workshop on the potential risks of antibody-dependent enhancement in human HIV vaccine trials*. Aids Research and Human Retroviruses, 1993. 9(12): p. 1175-84.
- 20 42. Szabo, J., *et al.*, *Strong correlation between the complement-mediated antibody-dependent enhancement of HIV-1 infection and plasma viral load*. Aids, 1999. 13(14): p. 1841-9.
43. Fust, G., *Enhancing antibodies in HIV infection*. Parasitology, 1997. 115 Suppl: p. S127-40.
- 25

Claims

1. An antibody or fragment thereof which is reactive with HIV-1 neutralizing antibody 2F5 (ECACC Accession No. 90091704) and which
5 inhibits or prevents the HIV-1 neutralization activity of antibody 2F5 and/or the binding of antibody 2F5 to gp41 of HIV-1.
2. The antibody or fragment thereof according to claim 1, which inhibits or prevents binding of antibody 2F5 to proteins or peptides that comprise the
10 amino acid sequence SEQ ID NO.1 (ELDKWA) or a variant thereof selected from the group consisting of SEQ ID NO.2 to SEQ ID NO 12.
3. The antibody or fragment thereof according to claim 1 or 2, which upon administration to a mammal elicits Ab3-type antibodies that have HIV-1
15 neutralizing activity, and/or that competes with antibody 2F5 for binding to proteins or peptides that comprise the amino acid sequence SEQ ID NO.1 (ELDKWA) or a variant thereof selected from the group consisting of SEQ ID NO.2 to SEQ ID NO 12.
- 20 4. The antibody or fragment thereof according to any one of claims 1 to 3, which is associated, preferably coupled, linked, or fused, with an immunogenic molecule, more preferably with a molecule that increases or intensifies a B cell response, and most preferably with a cytokine such as interleukin-4 (IL-4).
- 25 5. The antibody or fragment thereof according to any one of claims 1 to 4, wherein the antibody is an Ab2-type antibody which is anti-idiotypic to antibody 2F5.
- 30 6. The antibody or fragment thereof according to any one of claims 1 to 5, wherein the antibody is monoclonal and is produced by hybridoma cell line 3H6 (ECACC Acc. No. 01100279).

7. The antibody or fragment thereof according to any one of claims 1 to 6, in humanized form.
8. The antibody or fragment thereof according to any one of claims 1 to 7, for use as a screening tool or a diagnostic or therapeutic agent.
9. A hybridoma cell line producing an antibody according to any one of claims 1 to 6.
10. The hybridoma cell line according to claim 9, which is cell line 3H6 (deposited at PHLS, Porton Down, Salisbury, UK, under ECACC Acc. No. 01100279).
11. Use of an antibody or fragment thereof defined in any one of claims 1 to 8 for the manufacture of a pharmaceutical composition, particularly a vaccine, for prophylactic or therapeutic application against HIV-1 infection.
12. A pharmaceutical composition, particularly a vaccine, comprising an antibody or antibody fragment defined in any one of claims 1 to 8, and a pharmaceutically acceptable carrier.
13. An Ab3-type antibody or fragment thereof which is reactive with an antibody or antibody fragment according to any one of claims 1 to 8 and which competes with antibody 2F5 for binding to gp41 of HIV1 or to proteins or peptides that comprise the amino acid sequence SEQ ID NO.1 (ELDKWA) or a variant thereof selected from the group consisting of SEQ ID NO.2 to SEQ ID NO 12.
14. The Ab3-type antibody or antibody fragment according to claim 13 which neutralizes HIV-1.

- 22 -

15. The Ab3-type antibody or fragment thereof according to claim 13 or 14 for use as a screening tool or a diagnostic or therapeutic agent.

16. A pharmaceutical composition comprising an Ab3-type antibody or
5 antibody fragment according to anyone of claims 13 to 15, and a pharmaceutically acceptable carrier.

- 23 -

Abstract

The invention relates to Ab2-type anti-idiotypic antibodies and fragments thereof which mimic HIV-1 epitopes that are otherwise cryptic to the immune system and which antibodies or fragments thereof are directed against 5 potentially neutralizing anti-HIV-1 antibodies. The invention further relates to a hybridoma cell line 3H6 expressing the anti-idiotypic antibody and to pharmaceutical compositions containing said antibody or fragment thereof. The invention also relates to HIV-1 neutralizing Ab3-type antibodies elicited 10 upon administration of a said Ab2-type anti-idiotypic antibody or fragment thereof and to pharmaceutical compositions containing them. The invention also relates to the use of the present antibodies or fragments thereof as screening tools or as diagnostic or therapeutic agents.

1/3




FIG. 1




FIG. 2

2/3




FIG. 3




FIG. 4

3/3




FIG. 5

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.